


Amendments to the Claims

This listing of claims will replace all prior versions and listings of claims in the application:

Claims 1-65 are cancelled.

66. (currently amended) A compound having the structure:



or any pharmaceutically-acceptable salt thereof, wherein:

n is 1, 2 or 3 and o is 1, 2 or 3; wherein $n+o = 4$ or 5 and when n is 1 and Q^1 and Q^2 are both N, then both R^2 groups together are oxo, and when o is 1 and Q^1 and Q^2 are both N, then both $R^{2'}$ groups together are oxo;

m is independently at each instance 0, 1 or 2;

Q^1 is N or $C(R^2)$;

Q^2 is N or $C(R^2)$; wherein at least one of Q^1 and Q^2 is N;

Q^3 is $C(R^5)$;

Q^4 is $C(R^6)$;

Q^5 is $C(R^6)$;

Q^6 is $C(R^5)$;

R^1 is H or $-(C(R^2)(R^2))_m-R^g$;

R^2 is, independently, in each instance, H, C_{1-8} alkyl, C_{1-4} haloalkyl, $-O(C_{1-7}$ alkyl), $-N(C_{1-7}$ alkyl) R^a , or a C_{1-6} alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, $-OR^a$, $-OC(=O)R^b$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-NR^aC_{2-6}$ alkyl NR^aR^a and $-NR^aC_{2-6}$ alkyl OR^a ; wherein any two geminal R^2 groups may additionally be oxo;

$R^{2'}$ is, independently, in each instance, H, C_{1-8} alkyl, C_{1-4} haloalkyl, $-O(C_{1-7}$ alkyl), $-N(C_{1-7}$ alkyl) R^a , or a C_{1-6} alkyl substituted by 1, 2 or 3 substituents selected from halo, cyano, $-OR^a$, $-OC(=O)R^b$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$,

$-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$;
wherein any two geminal R^2 groups may additionally be oxo;

R^3 is, independently, in each instance, H, $C_{1-8}alkyl$, $C_{1-4}haloalkyl$,
 $-O(C_{1-7}alkyl)$, $-N(C_{1-7}alkyl)R^a$, or a $C_{1-6}alkyl$ substituted by 0, 1, 2 or 3 substituents
selected from halo, cyano, $-OR^a$, $-OC(=O)R^b$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$,
 $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$,
 $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$;
wherein any two geminal R^3 groups may additionally be oxo;

R^4 is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4
atoms selected from N, O and S, substituted by 0, 1, 2, 3 or 4 substituents selected
from $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$,
 $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$,
 $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$,
 $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$,
 $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$,
 $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$,
 $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$ and R^e ;

R^5 is independently, at each instance, H, $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano,
 $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$,
 $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$,
 $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$,
 $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$,
 $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$,
 $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$, $C_{1-3}alkylR^c$, $C_{1-3}alkylR^f$
and R^e ; or R^5 is a saturated or unsaturated 5-, 6- or 7-membered monocyclic or 6-, 7-,
8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from
N, O and S, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo
groups and the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from
 $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$,
 $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$,
 $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$,
 $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$,
 $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$,

$-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$ and R^c ;

R^6 is independently, at each instance, H, $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OH$, $-OC_{2-6}alkyl$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^b$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$ and R^c ;

R^a is independently, at each instance, H or R^b ;

R^b is independently, at each instance, phenyl, benzyl or $C_{1-6}alkyl$, the phenyl, benzyl and $C_{1-6}alkyl$ being substituted by 0, 1, 2 or 3 substituents selected from halo, $C_{1-4}alkyl$, $C_{1-3}haloalkyl$, $-OC_{1-4}alkyl$, $-NH_2$, $-NHC_{1-4}alkyl$, $-N(C_{1-4}alkyl)C_{1-4}alkyl$;

R^c is independently at each instance a saturated or unsaturated 5-, 6- or 7-membered monocyclic or 6-, 7-, 8-, 9-, 10- or 11-membered bicyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo groups;

R^d is independently at each instance $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$ or $-NR^aC_{2-6}alkylOR^a$;


R^e is independently at each instance $C_{1-6}alkyl$ substituted by 1, 2 or 3 substituents independently selected from R^d ;

R^f is independently at each instance R^c substituted by 1, 2 or 3 substituents independently selected from R^d ; and

R^g is independently at each instance a saturated or unsaturated 5-, 6- or 7-membered monocyclic or 6-, 7-, 8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo groups and the ring is substituted by 0, 1, 2 or 3 substituents selected from $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$,

-C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a,
-OC(=O)N(R^a)S(=O)₂R^b, -OC₂₋₆alkylNR^aR^a, -OC₂₋₆alkylOR^a, -SR^a, -S(=O)R^b,
-S(=O)₂R^b, -S(=O)₂NR^aR^a, -S(=O)₂N(R^a)C(=O)R^b, -S(=O)₂N(R^a)C(=O)OR^b,
-S(=O)₂N(R^a)C(=O)NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b,
-N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a,
-NR^aC₂₋₆alkylNR^aR^a and -NR^aC₂₋₆alkylOR^a.

67. (previously presented) A compound having the structure:



or any pharmaceutically-acceptable salt thereof, wherein:

n is 2 or 3;

m is independently at each instance 0, 1 or 2;

Q¹ is N or C(R²);

Q² is N or C(R²); wherein at least one of Q¹ and Q² is N;

Q³ is C(R⁵);

Q⁴ is C(R⁶);

Q⁵ is C(R⁶);

Q⁶ is C(R⁵);

R¹ is H or -(C(R²)(R²))_m-R^g;

R² is, independently, in each instance, H, C₁₋₈alkyl, C₁₋₄haloalkyl,

-O(C₁₋₇alkyl), -N(C₁₋₇alkyl)R^a, or a C₁₋₆alkyl substituted by 1, 2 or 3 substituents

selected from halo, cyano, -OR^a, -OC(=O)R^b, -SR^a, -S(=O)R^b, -S(=O)₂R^b,

-S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a,

-N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -NR^aC₂₋₆alkylNR^aR^a and -NR^aC₂₋₆alkylOR^a;

wherein any two geminal R² groups may additionally be oxo;

R³ is, independently, in each instance, H, C₁₋₈alkyl, C₁₋₄haloalkyl,

-O(C₁₋₇alkyl), -N(C₁₋₇alkyl)R^a, or a C₁₋₆alkyl substituted by 0, 1, 2 or 3 substituents

selected from halo, cyano, -OR^a, -OC(=O)R^b, -SR^a, -S(=O)R^b, -S(=O)₂R^b,

-S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a,

$-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$;
wherein any two geminal R^3 groups may additionally be oxo;

R^4 is an unsaturated 5- or 6-membered monocyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, substituted by 0, 1, 2, 3 or 4 substituents selected from $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$ and R^c ;

R^5 is independently, at each instance, H, $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^bR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$, $C_{1-3}alkylR^c$, $C_{1-3}alkylR^f$ and R^c ; or R^5 is a saturated or unsaturated 5-, 6- or 7-membered monocyclic or 6-, 7-, 8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo groups and the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$ and R^c ;

R^6 is independently, at each instance, H, $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OH$, $-OC_{2-6}alkyl$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$,

$-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^b$, $-N(R^a)C(=O)R^b$,
 $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$,
 $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$ and R^c ;

R^a is independently, at each instance, H or R^b ;

R^b is independently, at each instance, phenyl, benzyl or $C_{1-6}alkyl$, the phenyl, benzyl and $C_{1-6}alkyl$ being substituted by 0, 1, 2 or 3 substituents selected from halo, $C_{1-4}alkyl$, $C_{1-3}haloalkyl$, $-OC_{1-4}alkyl$, $-NH_2$, $-NHC_{1-4}alkyl$, $-N(C_{1-4}alkyl)C_{1-4}alkyl$;

R^c is independently at each instance a saturated or unsaturated 5-, 6- or 7-membered monocyclic or 6-, 7-, 8-, 9-, 10- or 11-membered bicyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo groups;

R^d is independently at each instance $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$ or $-NR^aC_{2-6}alkylOR^a$;

R^e is independently at each instance $C_{1-6}alkyl$ substituted by 1, 2 or 3 substituents independently selected from R^d ;

R^f is independently at each instance R^c substituted by 1, 2 or 3 substituents independently selected from R^d ; and

R^g is independently at each instance a saturated or unsaturated 5-, 6- or 7-membered monocyclic or 6-, 7-, 8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo groups and the ring is substituted by 0, 1, 2 or 3 substituents selected from $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_{2-6}alkylNR^aR^a$, $-OC_{2-6}alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$.

68. (previously presented) The compound according to Claim 67, wherein Q^1 is N.

69. (previously presented) The compound according to Claim 67, wherein Q^2 is N.

70. (previously presented) The compound according to Claim 67, wherein:
 Q^1 is N; and
 Q^2 is N.

71. (previously presented) The compound according to Claim 67, wherein R^1 is H.

72. (previously presented) The compound according to Claim 67, wherein R^1 is $-(C(R^2)(R^2))_m-R^g$.

73. (previously presented) The compound according to Claim 67, wherein R^2 is, in each instance, H.

74. (previously presented) The compound according to Claim 67, wherein at least one R^2 group is selected from C_{1-8} alkyl, C_{1-4} haloalkyl, $-O(C_{1-7}$ alkyl), $-N(C_{1-7}$ alkyl) R^a , oxo and C_{1-6} alkyl substituted by 0, 1, 2 or 3 substituents selected from halo, cyano, $-OR^a$, $-OC(=O)R^b$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-NR^aC_{2-6}$ alkyl NR^aR^a and $-NR^aC_{2-6}$ alkyl OR^a .

75. (previously presented) The compound according to Claim 67, wherein R^3 is, in each instance, H.

76. (previously presented) The compound according to Claim 67, wherein at least one R^3 group is selected from C_{1-8} alkyl, C_{1-4} haloalkyl, $-O(C_{1-7}$ alkyl), $-N(C_{1-7}$ alkyl) R^a , oxo and C_{1-6} alkyl substituted by 0, 1, 2 or 3 substituents selected

from halo, cyano, $-OR^a$, $-OC(=O)R^b$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$,
 $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$,
 $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-NR^aC_{2-6}alkylNR^aR^a$ and $-NR^aC_{2-6}alkylOR^a$.

77. (previously presented) The compound according to Claim 67, wherein R^4 is an unsaturated 6-membered monocyclic ring containing 1, 2 or 3 atoms selected from N, O and S, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo groups and the rings are substituted by 0, 1, 2, 3 or 4 substituents selected from $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_2-6alkylNR^aR^a$, $-OC_2-6alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$ and R^e .

78. (previously presented) The compound according to Claim 67, wherein R^4 is an unsaturated 5-membered monocyclic ring containing 1, 2 or 3 atoms selected from N, O and S, but no more than one N, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo groups and the rings are substituted by 0, 1, 2, 3 or 4 substituents selected from $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_2-6alkylNR^aR^a$, $-OC_2-6alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$, $-S(=O)_2N(R^a)C(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkylNR^aR^a$, $-NR^aC_{2-6}alkylOR^a$ and R^e .

79. (previously presented) The compound according to Claim 67, wherein R^5 is independently, at each instance, H, $C_{1-8}alkyl$, $C_{1-4}haloalkyl$, halo, cyano, $-C(=O)R^b$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC(=O)N(R^a)S(=O)_2R^b$, $-OC_2-6alkylNR^aR^a$, $-OC_2-6alkylOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-S(=O)_2N(R^a)C(=O)R^b$, $-S(=O)_2N(R^a)C(=O)OR^b$,

-S(=O)₂N(R^a)C(=O)NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b,
-N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a,
-NR^aC₂₋₆alkylNR^aR^a, -NR^aC₂₋₆alkylOR^a, C₁₋₃alkylR^c, C₁₋₃alkylR^f and R^e; or R⁵ is a
saturated or unsaturated 5-, 6- or 7-membered monocyclic or 6-, 7-, 8-, 9-, 10- or
11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S,
wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo groups and the
ring is substituted by 0, 1, 2, 3 or 4 substituents selected from C₁₋₈alkyl, C₁₋₄haloalkyl,
halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a,
-OC(=O)R^b, -OC(=O)NR^aR^a, -OC(=O)N(R^a)S(=O)₂R^b, -OC₂₋₆alkylNR^aR^a,
-OC₂₋₆alkylOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -S(=O)₂N(R^a)C(=O)R^b,
-S(=O)₂N(R^a)C(=O)OR^b, -S(=O)₂N(R^a)C(=O)NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b,
-N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b,
-N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkylNR^aR^a, -NR^aC₂₋₆alkylOR^a and R^e.

80. (previously presented) The compound according to Claim 67, wherein
R⁵ is, at each instance, H.

81. (previously presented) The compound according to Claim 67, wherein
at least one R⁵ is C₁₋₈alkyl, C₁₋₄haloalkyl, halo, cyano, -C(=O)R^b, -C(=NR^a)NR^aR^a,
-OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC(=O)N(R^a)S(=O)₂R^b, -OC₂₋₆alkylNR^aR^a,
-OC₂₋₆alkylOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -S(=O)₂N(R^a)C(=O)R^b,
-S(=O)₂N(R^a)C(=O)OR^b, -S(=O)₂N(R^a)C(=O)NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b,
-N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b,
-N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkylNR^aR^a, -NR^aC₂₋₆alkylOR^a, C₁₋₃alkylR^c, C₁₋₃alkylR^f
and R^e.

82. (previously presented) The compound according to Claim 67, wherein
at least one R⁵ is a saturated or unsaturated 5-, 6- or 7-membered monocyclic or 6-, 7-,
8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from
N, O and S, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo
groups and the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from
C₁₋₈alkyl, C₁₋₄haloalkyl, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,
-C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC(=O)N(R^a)S(=O)₂R^b,

-OC₂₋₆alkylNR^aR^a, -OC₂₋₆alkylOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a,
-S(=O)₂N(R^a)C(=O)R^b, -S(=O)₂N(R^a)C(=O)OR^b, -S(=O)₂N(R^a)C(=O)NR^aR^a, -NR^aR^a,
-N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
-N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkylNR^aR^a, -NR^aC₂₋₆alkylOR^a and
R^c.

83. (previously presented) The compound according to Claim 67, wherein at least one R⁵ is an unsaturated 5-, 6- or 7-membered monocyclic ring containing 0, 1, 2 or 3 atoms selected from N, O and S, wherein the carbon atoms of the ring are substituted by 0, 1 or 2 oxo groups and the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from C₁₋₈alkyl, C₁₋₄haloalkyl, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC(=O)N(R^a)S(=O)₂R^b, -OC₂₋₆alkylNR^aR^a, -OC₂₋₆alkylOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -S(=O)₂N(R^a)C(=O)R^b, -S(=O)₂N(R^a)C(=O)OR^b, -S(=O)₂N(R^a)C(=O)NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkylNR^aR^a, -NR^aC₂₋₆alkylOR^a and R^c.

84. (previously presented) The compound according to Claim 67, wherein at least one R⁵ is an unsaturated 6-membered monocyclic ring containing 0, 1 or 2 N atoms, wherein the ring is substituted by 0, 1, 2, 3 or 4 substituents selected from C₁₋₈alkyl, C₁₋₄haloalkyl, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC(=O)N(R^a)S(=O)₂R^b, -OC₂₋₆alkylNR^aR^a, -OC₂₋₆alkylOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -S(=O)₂N(R^a)C(=O)R^b, -S(=O)₂N(R^a)C(=O)OR^b, -S(=O)₂N(R^a)C(=O)NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkylNR^aR^a, -NR^aC₂₋₆alkylOR^a and R^c.

85. (previously presented) The compound according to Claim 67, wherein at least one R⁶ is selected from C₁₋₈alkyl, C₁₋₄haloalkyl, halo, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC(=O)N(R^a)S(=O)₂R^b, -OC₂₋₆alkylNR^aR^a, -OC₂₋₆alkylOR^a, -SR^a, -S(=O)R^b,

-S(=O)₂R^b, -S(=O)₂NR^aR^a, -S(=O)₂N(R^a)C(=O)R^b, -S(=O)₂N(R^a)C(=O)OR^b,
-S(=O)₂N(R^a)C(=O)NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a,
-N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkylNR^aR^a,
-NR^aC₂₋₆alkylOR^a and R^c.